









WHAT IS CLAIMED IS:


1. A method for treating ocular hypertension or glaucoma which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of Table 1:
- 5
- Table 1














emindole DA




epi-emindole DA










emindole PA




emindole DB












emindole SB














Fumitremorgen A




Janthitrem B




Janthitrem C




Janthitrem E




Janthitrem F




Janthitrem G




Lolitre<sup>m</sup> A




Lolitre<sup>m</sup> B




Lolitre<sup>m</sup> C








Lolitre<sup>m</sup> E







Lolitre<sup>m</sup> F



Lolitre<sup>m</sup> H





or a pharmaceutically acceptable salt, enantiomer, diastereomer, tautomer or mixture thereof.

2. The method according to Claim 1 wherein the compound of formula I is applied as a topical formulation.

3. The method according to claim 3 wherein the topical  
5 formulation is a solution or suspension.

4. The method of Claim 3, which comprises administering a second active ingredient, concurrently or consecutively, wherein the second active ingredient is a hypotensive agent selected from a  $\beta$ -adrenergic blocking agent,  
10 adrenergic agonist, a parasympathomimetic agent, a carbonic anhydrase inhibitor, EP4 agonist and a prostaglandin or a prostaglandin derivative.

5. The method according to claim 4 wherein the  $\beta$ -adrenergic blocking agent is timolol, levobunolol, carteolol, optipranolol, metapranolol or  
15 betaxolol; the parasympathomimetic agent is pilocarpine, carbachol, or phospholine iodide; adrenergic agonist is iopidine, brimonidine, epinephrine, or dipivephrin, the carbonic anhydrase inhibitor is dorzolamide, acetazolamide, metazolamide or brinzolamide; the prostaglandin is latanoprost or rescula, and the prostaglandin derivative is a hypotensive lipid derived from PGF2 $\alpha$  prostaglandins.

20 6. A method according to claim 2 in which the topical formulation contains xanthan gum or gellan gum.

7. A method for treating macular edema, macular degeneration,  
25 for providing a neuroprotective effect, increasing retinal and optic nerve head blood velocity or increasing retinal and optic nerve oxygen tension which comprises administering to a patient in need of such treatment a pharmaceutically effective amount of a compound as recited in claim 1

8. The method according to Claim 7 wherein the compound of formula I is applied as a topical formulation in the form of a solution or suspension.

9. The method of Claim 8, which comprises administering a second active ingredient, concurrently or consecutively, wherein the second active  
35 ingredient is a hypotensive agent selected from a  $\beta$ -adrenergic blocking agent,

adrenergic agonist, a parasympathomimetic agent, a carbonic anhydrase inhibitor, EP4 agonist and a prostaglandin or a prostaglandin derivative.

10. The method according to claim 9 wherein the  $\beta$ -adrenergic  
5 blocking agent is timolol, levobunolol, carteolol, optipranolol, metapranolol or  
betaxolol; the parasympathomimetic agent is pilocarpine, carbachol, or phospholine  
iodide; adrenergic agonist is iopidine, brimonidine, epinephrine, or dipivephrin, the  
carbonic anhydrase inhibitor is dorzolamide, acetazolamide, metazolamide or  
brinzolamide; the prostaglandin is latanoprost or rescula, and the prostaglandin  
10 derivative is a hypotensive lipid derived from PGF2 $\alpha$  prostaglandins.

11. A method according to claim 8 in which the topical formulation  
contains xanthan gum or gellan gum.